

# Lecture 12.5

Basics of Sets and Functions

# What's a Set?

**Defn:** A **set** is an unordered collection of **distinct** objects, called **elements** or members of the set.

$a \in A$  denotes that  $a$  is a member of  $A$ , and  $a \notin A$  denotes that  $a$  is not a member of  $A$ .

## Examples:

$V = \{a, e, i, o, u\}$ , is the set of vowels in English alphabet.

$E = \{2, 4, 6, 8, 10\}$ , is the set of positive even integers  $\leq 10$ .

$E = \{x \mid x \text{ is a positive even integer } \leq 10\}$

$\mathbb{Q}^+ = \{x \in \mathbb{R} \mid x = \frac{p}{q}, \text{ for some positive integers } p \text{ and } q\}$

*Roster notation.*

*Set builder notation.*

**Note:** It is not necessary that members of a set should have a common property.

For instance,  $\{99, \text{Bob}, \text{Jupiter}\}$  is a valid set.

# More about Sets

**Definition:** Two sets are **equal** if and only if they have the **same elements**. In other words, if  $A$  and  $B$  are sets, then  $A = B$  if and only if  $\forall x(x \in A \iff x \in B)$ .

**Example:**  $\{1,2,3\} = \{1,3,2\}$  because they contain the same elements and the order does not matter. It also does not matter whether one element is listed more than once, therefore,  $\{1,2,3,3,2,2\} = \{1,2,3\}$ .

**Definition:** A set that contains no elements is called the **empty set** and denoted by  $\emptyset$ .

A set with just one element is called a **singleton set**.

**Note:** Do not confuse  $\emptyset$  with  $\{\emptyset\}$ ,  $\emptyset$  is the empty set and  $\{\emptyset\}$  is a singleton set.

# Subsets and Cardinality

**Defn:** The set  $A$  is a **subset** of  $B$  iff every element of  $A$  is also an element of  $B$ .

$A \subseteq B$  denotes that  $A$  is a subset of  $B$ .

**Proving  $A \subseteq B, A \not\subseteq B, A = B$ :**

- To prove  $A \subseteq B$ , show that if  $x \in A$ , then  $x \in B$ .
- To prove  $A \not\subseteq B$ , find an  $x$  in  $A$  such that  $x \notin B$ .
- To prove  $A = B$ , show that  $A \subseteq B$  and  $B \subseteq A$ .

**Defn:** Let  $S$  be a set. If there are exactly  $n$  distinct elements in  $S$ , where  $n$  is a nonnegative integer, we say that  $S$  is a finite set and that  $n$  is the **cardinality** of  $S$ . The cardinality of  $S$  is denoted by  $|S|$ . A set is said to be **infinite** if it is not finite.

# Ordered Tuple

**Defn:** The **ordered  $n$ -tuple**  $(a_1, a_2, \dots, a_n)$  is the ordered collection that has  $a_1$  as its first element,  $a_2$  as its second element,  $\dots$ , and  $a_n$  as its  $n$ th element. Ordered 2-tuples are called **ordered pairs**.

Two ordered  $n$ -tuples, say  $(a_1, a_2, \dots, a_n)$  and  $(b_1, b_2, \dots, b_n)$ , are **equal** if and only if each corresponding pair of their elements are equal, i.e.,  $a_i = b_i$  for  $i = 1, 2, \dots, n$ .

# Cartesian Product

**Defn:** Let  $A$  and  $B$  be sets. The **Cartesian product** of  $A$  and  $B$ , denoted by  $A \times B$ , is the set of all ordered pairs  $(a, b)$ , where  $a \in A$  and  $b \in B$ . Hence,

$$A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$$

**Example:** Let  $A = \{a, b\}$  and  $B = \{1, 2, 3\}$ . Then

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

$$B \times A = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$$

**Definition:** The cartesian product of the sets  $A_1, A_2, \dots, A_n$ , denoted by  $A_1 \times A_2 \times \dots \times A_n$ , is the set of ordered  $n$ -tuples  $(a_1, a_2, \dots, a_n)$ , where  $a_i$  belongs to  $A_i$ , for  $i = 1, 2, \dots, n$ .

# Set Operations

Let  $A$  and  $B$  be two sets. Then the following operation can be defined on them,

**Union:** Denoted by  $A \cup B$ , is the set of all the elements that are either in  $A$  or  $B$ , or in both.

**Intersection:** Denoted by  $A \cap B$ , is the set of all the elements that are in both  $A$  and  $B$ .

$A$  and  $B$  are **disjoint**, if  $A \cap B = \emptyset$ .

**Difference:** Denoted by  $A - B$ , is the set of all the elements that are in  $A$  but not in  $B$ .

**Complement:** Let  $U$  be the universal set. The complement of the set  $A$ , denoted by  $\bar{A}$ , is the complement of  $A$  with respect to  $U$ , i.e.,  $U - A$ .

**Note:** Union and intersection of more than two sets defined as the natural extension of union and intersection of two sets.



# Set Identities

**Identity Laws:**  $A \cap U = A$   
 $A \cup \emptyset = A$

**Domination Laws:**  $A \cup U = U$   
 $A \cap \emptyset = \emptyset$

**Idempotent Laws:**  $A \cup A = A$   
 $A \cap A = A$

**Absorption Laws:**  $A \cup (A \cap B) = A$   
 $A \cap (A \cup B) = A$

**Complementation Law:**  $\overline{\overline{A}} = A$

**De Morgan's Laws:**  $\overline{A \cap B} = \overline{A} \cup \overline{B}$   
 $\overline{A \cup B} = \overline{A} \cap \overline{B}$

**Complement Laws:**  $A \cup (A \cap B) = A$   
 $A \cap (A \cup B) = A$

**Commutative Laws:**  $A \cup B = B \cup A$   
 $A \cap B = B \cap A$


**Associative Laws:**  $A \cup (B \cup C) = (A \cup B) \cup C$   
 $A \cap (B \cap C) = (A \cap B) \cap C$

**Distributive Laws:**  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$   
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$




# Functions

**Defn:** Let  $A$  and  $B$  be nonempty sets. A **function**  $f$  from  $A$  to  $B$  is an **assignment** of **exactly one element** of  $B$  to each element of  $A$ . We write  $f(a) = b$  if  $b$  is the unique element of  $B$  assigned by the function  $f$  to the element  $a$  of  $A$ . If  $f$  is a function from  $A$  to  $B$ , we write  $f: A \rightarrow B$ .  $A$  is called the **domain** of  $f$  and  $B$  is called the **image** or **range** of  $f$ .




*A function*




*Not a function*

# One-to-One Functions

**Defn:** A function  $f$  is said to be **one-to-one** or an **injection**, if and only if  $f(a) = f(b)$  implies that  $a = b$  for all  $a$  and  $b$  in the domain of  $f$ .




*An injection*




*Not an injection*

# Onto Functions

**Definition:** A function  $f$  is said to be **onto** or a **surjection**, if and only if for every element  $b \in B$  there is an element  $a \in A$  with  $f(a) = b$ .




*A surjection*




*Not a surjection*

# Bijjective Functions

**Defn:** A function  $f$  is said to be a **bijection**, if and only if it is both **one-to-one** and **onto**.



*A bijection*



*Not a bijection*

# Inverse Function and Composition of Functions

**Defn:** Let  $f$  be a bijection from  $A$  to  $B$ . The **inverse function** of  $f$ , denoted by  $f^{-1}$ , is the function that assigns to an element  $b$  of  $B$  the unique element  $a$  in  $A$  such that  $f(a) = b$ , i.e.,  $f^{-1}(b) = a$ .

**Defn:** Let  $g$  be a function from  $A$  to  $B$  and let  $f$  be a function from  $B$  to  $C$ . The **composition** of the functions  $f$  and  $g$ , denoted by  $f \circ g$  for all  $a \in A$ , is defined by

$$(f \circ g)(a) = f(g(a))$$